



## **WART INSTRUCTIONS**

### **WART HELPFUL HINTS:**

If your wart has been treated with liquid nitrogen it may blister, turn red, and even scab over. This is normal. Wait for these symptoms to subside before further treatment with topical medications. If you get a blister that is uncomfortable you can poke the blister with a sterile needle to let the fluid out. Otherwise, if it is not bothersome leave the treated area alone and the blister will resolve.

### **PRESCRIPTION WART PEEL INSTRUCTIONS:**

Wait until the wart has “healed” from the liquid nitrogen treatment, then begin topical treatment. Apply wart treatment each night & cover with duct tape. Remove the tape in the AM. Twice weekly file the wart with a disposable emery board (dispose of it after each use).

**STOP** topical treatments if you notice significant irritation, redness, or ulceration and allow the area to heal.

### **OVER THE COUNTER WART PREPARATIONS:**

Use as instructed by the directions included on the purchased product. Use until the wart has resolved. If significant irritation develops, hold treatment until the irritation resolves.

- Dr. Scholl’s “Clear Away” with 40% salicylic acid
- Duofilm
- Compound W one step pads
- Compound W Wart Remover gel with 17% salicylic acid

**Please contact the office if you have any questions or concerns**



# **Cosmetic SK's, Skin Tags, Hyfreaction, and Small Angioma Removals:**

1:\$50

2-15:\$150

16-30:\$290

**\*\*It will be up to the provider if it is considered cosmetic  
once they are evaluated in the exam room.\*\***